




Know the FACTS. Understand the RISKS.

THE FACTS



people in the U.S. live with diabetes.¹



Nearly 1 in 4 adults are living with diabetes.¹



of patients with diabetes may experience a foot ulcer during their lifetime.²



of people with a **healed diabetic foot ulcer** will develop a new ulcer within a year.²

THE RISKS



About 60% of people living with diabetes will develop neuropathy, eventually leading to a foot ulcer.³



People with diabetes are 2-4 times more likely to have a cardiovascular disease.⁴




Every 30 seconds in the world, a lower limb is amputated due to diabetes.⁵



After amputation, 55% require an amputation on the second leg within 2-3 years.⁶

TREATMENT



Perform foot examinations on your patients checking for neuropathy, infection, wounds, ulcers, foot swelling or burning sensations that may be present.



Amputation risk is reduced by almost 50% when care is provided by a multidisciplinary wound care team.⁴

Advanced wound care treatments include **debridement, total contact casting or offloading, new tissue growth, negative pressure wound therapy and hyperbaric oxygen therapy (HBO).**

Our center provides advanced wound care treatments with the knowledge and resources to help treat your patients' wounds. **Come Heal With Us! Learn more at [Healogics.com](https://healogics.com).**



FIND. TREAT. HEAL.™

1.CDC; 2.Armstrong, Boulton, Bus, New England Journal of Medicine, 2017; 3.Oliver TI, Mutluoglu M. Diabetic Foot Ulcer; 4.AHA, 5.International Diabetes Federation and International Working Group of the Diabetic Foot . In: Time to act. Bakker K, Foster AV; 6.Pandian G, Hamid F, Hammond M. Rehabilitation of the Patient with Peripheral Vascular Disease and Diabetic Foot Problems.
©2024 by Healogics, LLC. All Rights Reserved.