


Know the **FACTS**. Understand the **RISKS**.

THE FACTS



people in the U.S. live
with diabetes.¹



Nearly 1 in 4 adults are living
with diabetes.¹

15-25%

of patients with diabetes may
experience a foot ulcer during
their lifetime.²

40%

of people with a **healed
diabetic foot ulcer** will develop a
new ulcer within a year.²

THE RISKS

60%

About 60% of people living
with diabetes will develop
neuropathy, eventually
leading to a foot ulcer.³

2-4x

People with diabetes
are 2-4 times more
likely to have a
cardiovascular disease.⁴



Every 30 seconds
in the world, a lower limb is
amputated due to diabetes.⁵

55%

After amputation, 55% require
an amputation on the second leg
within 2-3 years.⁶

TREATMENT



**Perform foot
examinations** on your
patients checking for
neuropathy, infection,
wounds, ulcers, foot
swelling or burning
sensations that may
be present.



Amputation risk is
reduced by almost 50%
when care is provided by a
multidisciplinary wound
care team.⁴

Advanced wound care
treatments include
**debridement, total contact
casting or offloading, new tissue
growth, negative pressure
wound therapy and hyperbaric
oxygen therapy (HBO).**

Our center provides advanced wound care treatments with the knowledge and resources to help treat your patients' wounds.
Come Heal With Us! Learn more at [Healogics.com](https://www.healogics.com).

Healogics®

FIND. TREAT. HEAL.™